

Effect of some abiotic factors on the growth of *Oreochromis niloticus* Linnaeus, 1758 (Perciformes: Cichlidae) in tropical ecosystems (Cameroon)

Authors:

Mohamed Nsangou
Nchoutndignigni^{1,2*},
Sévilor Kekeunou²,
Gidéon Ajeegah³,
Benoit Bapfubusa²,
Abraham Fomena⁴

Institution:

1 Laboratory of Zoology, Higher
Teacher Training College,
University of Yaoundé I, Cameroon

2 Laboratory of Zoology, Faculty of
Science, University of Yaoundé I,
Cameroon

3 Laboratory of Hydrobiology and
Environment, Faculty of Science,
University of Yaoundé I, Cameroon

4 Laboratory of Parasitology and
Ecology, Faculty of Science,
University of Yaoundé I, Cameroon

Corresponding author:

Mohamed Nsangou
Nchoutndignigni

ABSTRACT:

The Growth of *Oreochromis niloticus* (2.6g initial average size) was studied in 100m² earthen ponds for 180 days under tropical field environment. Varying fertilizing regimes consisting of *Pennisetum purpureum* (PP) and *Musa sapientum* (MS), two locally available weedy grasses were applied in duplicates in the experimental ponds, at the rate of 0.1-0.2 kg dry matter per m² per day. Water quality parameters were monitored bimonthly between 8:00am-9:00am . Final average fish weight varied significantly ($p < 0.05$) as follows: 43.85g, 35.5g, 59.5g, 24.9g and 50g, 2 respectively for PP, MS, MS+PP, T₀ and T₁. Corresponding daily growth were 2.2g/d, 0.1g/d, 0.3g/d, 0.1g/d and 0.2g/d. Survival rate equally varied significantly as follow: 60 %, 67 %, and 98 %, 70 % and 97 % respectively. During the study period, the results indicated the water quality were permissible limits and can be used for livestock of *Oreochromis niloticus* .

Keywords:

Oreochromis niloticus, fish farming, pond, water quality, fertilization.