

Original Research

Malacofaunal distribution, abundance and diversity of the Nakana Lake, Dhule (MS) India

Authors:

Patil MU¹
Patole SS²,
Bhoi SS³ and
Ahirrao KD⁴

Institution:

1. Dept. of Zoology, V.U. Patil Arts, and Late Dr. B. S. Desale science college, Sakri, Dist-Dhule 424304.

2. Dept. of Zoology, V,V,M.'s S.G. Patil Arts Science and Commerce College, Sakri, District-Dhule.

3. Dept. of Zoology, Uttamrao. Patil Arts, and science college, Dahivel, Tal. Sakri, Dist-Dhule.

4. Head, Dept of Zoology, Rani Laxmibai College, Parola, Dist-Jalgaon 425 111.

Corresponding author:

Patil Manisha U

ABSTRACT:

Freshwater Molluscs distribution and diversity status of Nakana Lake, Dhule, were studied during June 2013 to May 2014. The samples were collected at every fortnight from two sites. A total of 13 species were reported, out of which 10 species were Gastropods and three species were Pelecypods; those belonged to four orders, 10 families and 12 genera. The abundance of molluscs was also studied. The average number of animals were indicated per season: i.e. monsoon, winter and summer. Maximum species were collected in the summer season at both the sites. Molluscan diversity were calculated by the various diversity indices such as Shannon-Wiener diversity index (H), Simpson's Dominance index (D), Simpson's index of diversity (I-D), Simpson's Evenness(E) and Pielou's Evenness index (J).

Keywords:

Molluscan diversity, Seasonal variations, Abundance, Biological Indices.

Email Id:

rajmany155@gmail.com

Article Citation:

Patil MU, Patole SS, Bhoi SS and Ahirrao KD

Malacofaunal distribution, abundance and diversity of the Nakana Lake, Dhule (MS) India.

Journal of Research in Biology (2016) 6(2): 1977-1983

Dates:

Received: 09 Dec 2016

Accepted: 18 Jan 2016

Published: 04 March 2016

Web Address:

<http://jresearchbiology.com/documents/RA0584.pdf>

This article is governed by the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which gives permission for unrestricted use, non-commercial, distribution and reproduction in all medium, provided the original work is properly cited.

INTRODUCTION

The mollusca are soft bodied animals. They move on the land and still dependent on a moist ground, in an excess of cold or hot and dry climate they enter in the state of dormancy. The Phylum Mollusca are classified into seven classes viz. Aplacophora, Polyplacophora, Monoplacophora, Gastropoda, Scaphopoda, Pelecypoda, and Cephalopoda. Gastropoda is the largest class more than 80,000 species are recorded throughout the world. (Hyman, 1967).

Mollusca prove immensely beneficial in economic and medicinal values. Like all animals, the molluscs are not considered by men to be either useful or destructive. Gastropod plays an important role in balancing the nature. The shells were used in Unani and Ayurvedic medicine system for meditation, in the production of humus, in the control of fungi, algae, lichens and also as predators and parasites; but in nature the molluscs are hunted and eaten by predators (Ahirrao, 2003).

In scientific studies, molluscs prove key role in study of drug action on heart, hormones, enzymes and antitoxins-especially in relation to immunological haematology. These are suitable bioindicators for some

radio-active and chemical pollution in the coastal areas of water bodies. The snails and slugs are important to man because of the damage they do in agriculture, horticulture and forestry. Furthermore, they are of importance in medical and veterinary practice, since they serve as intermediate host for certain parasitic worms of man and domestic animals viz. Schistosome parasites (Ahirrao, 2003; Ahirrao and Khedkar, 2012).

MATERIALS AND METHODS

Geographical area of Nakana Lake

The present study area is selected from one of the man-made Lake situated on the Panzara River, 6 km away from Dhule city. It runs parallel to the National highway number: 6 (NH-6). Geographical location of the lake is as follows; Longitudes:- 74.7417477 and Latitudes:- 20.9095349. It is an earthen lake having catchment area of 945 m³, with a maximum height is 18.41 m. It also receives water from Haranmall Lake. The lake water is used for drinking, irrigation and aquaculture purpose. One side of it is covered by a well developed forest with grass trees, herbs and shrubs. The grass starts in the forest drying from the start of post monsoon, frequently cattle grazing is observed in this



Figure 1. Map Showing Study area, Nakana Lake, Dhule (MS) India




Figure 2. Photographs showing the study areas (i) S-I and (ii) S-II.

area. During present investigation two sites were selected for the study namely S-I and S-II.

Sampling and preservation

The Molluscan species were collected by hand picking method for the big specimens and for the smallest one, a sieve for soil samples were used. All specimens were brought to the laboratory, washed thoroughly, and preserved in 4 % formalin. The shells were dried at room temperature, and then separated in polythene bags. Identification was done on the basis of Preston (1915) Rao (1989) and Tonapi (1980). The unidentified molluscan shells are identified and classified upto the generic and species level by Zoological survey of India, Pune, Maharashtra State, India. Data were collected fortnightly, pooled seasonally viz. monsoon,

winter and summer from June 2013 to May 2014.

Data analysis

Quantitative and qualitative estimation of molluscan fauna were carried out using quadrat method (Micheal, 1984, Subba Rao, 1989). At least four quadrat (1 X 1m) were sampled from each site and the average of these were considered as density and species richness per m² for that area.

Formulae for indices

Total number of species, total number of individual in a sample and different diversity indices. From the present data were calculated using the following indices Shannon-Weiner Index (H) (1949), Simpson’s Dominance Index (D), Simpson (1949), Simpson’s Index of diversity (1-D), Simpson’s reciprocal

Table 1. The Molluscan species richness, abundance and diversity indices of Nakana Lake and Dhule are tabulated.

S. No.	Index	Site -I			Site-II		
		M	W	S	M	W	S
1	Species Richness (S)	13	13	13	12	12	12
2	Species abundance(N)	54	80	137	37	50	104
3	Shannon-Weiner Index(H)	3.8134	3.0866	6.5137	3.058	2.2611	5.2641
4	Simpson’s Dominance Index (D)	0.0887	0.1183	0.1212	0.1216	0.1461	0.1376
5	Simpson’s Index of Diversity (1-D)	0.9113	0.8817	0.8788	0.8784	0.8539	0.8624
6	Simpson’s reciprocal Index (1/D)	11.273	8.453	8.2508	8.2226	6.8446	7.2674
7	Simpson’s Evenness (E)	0.8671	0.6502	0.6346	0.6853	0.5703	0.6056
8	Pielou’s Evenness (J)	1.4867	1.2033	2.5395	1.2306	0.9099	2.1184




Figure 3. Pie Charts showing percentage of annual distribution of species at two sites: S-I and S-II respectively

Index (1/D), Simpson's Evenness (E) and Pielou's evenness Index (J) Pielou(1975). The following index values were used for calculation.

- 1.Shannon – Weiner Index (H): $H = -\sum P_i (\ln P_i)$,
- 2.Simpson's Dominance Index (D): $D = \sum n(n-1)/N(N-1)$,
- 3.Simpson's Index of Diversity (D_1): $D_1 = 1-D$,
- 4.Simpson's reciprocal Index (D_2): $D_2 = 1/D$,
- 5.Simpson's Evenness (E): $E = D_2/S$
- 6.Pielou's evenness Index (J): $J = H/\ln^*S$

Where,

S = Number of species.

N = Total number of individual of all species.

$P_i = A/T$ where A is number of each species in the sample,

T = Total number of individual of all species in the sample.

n = Total number of individuals of particular species.

RESULTS AND DISCUSSION

Seasonal variations

In the present investigation 462, samples were collected during June 2013 to May 2014 from the both sites i.e. Site I (Nakana Lake) and site II (Dhule). Totally, 13 species were recorded, which belongs to 4 orders, 10 families and 12 genera. Among them

Gastropods found to be dominant over Pelecypods, it contributes 86.14% and 13.65 % respectively. Sharma *et.al.* (2009), recorded similar predominance of Gastropoda over Pelecypoda. From Site I =13 species, and from Site II =12 species were recorded. This is shown in Fig.3.

Density of Mollusca at Site-I ranges between 54 organism/ m^2 in monsoon , 80 org./ m^2 in winter and 137 org./ m^2 in summer season, whereas at Site-II, least 37 org./ m^2 in Monsoon, moderate number 50 org./ m^2 in Winter and maximum i.e. 104 org./ m^2 in Summer.

Observations showed that highest population of Molluscs were recorded during summer from both the sites that are corresponding with Hussein *et al.*,(2011). He reported March-May pick in his study. The snail abundance was maximum during the summer which might be due to ecological conditions like warm temperature, activates the process of decomposition. A large amount of decomposers is responsible for the deposition of organic matter and macrophytes at the bottom of the lake.

Total decline in molluscan population was




Figure 4. The column chart shows comparison among different diversity indices of recorded molluscan species, at Nakana Lake, Dhule.

observed at monsoon season due to self dilution of the body fluid. The sensitive molluscs could not adjust the fluctuating osmotic balance suddenly hence the mortality was very high during the period of monsoon season. The salinity and temperature dropped down which is made the condition adverse for the animal because of heavy rainfall, Patole (2010).

In winter at both sites population of molluscs found to be moderate in number, because of the suitable ecosystem but molluscs are poikilotherms, and they cannot tolerate an excess cold environment and hence they may enter in to the state of hibernation. On the other hand El-Kady *et al.*, (2000) was recorded low number of snails during winter (Jan and Feb) in Sinai Peninsula.

Biological indices

The diversity of molluscs mainly depends up on biotic, abiotic factors and stability of the ecosystem. Molluscan diversity is calculated by the diversity status, richness, abundance, habitat and various diversity indices. Values are shown in table no. 1.

For proper understanding the Biotic community, it is essential to work out some indices Sharma *et al.*,

(2010). There has been quite large fluctuation between the level of abundance and showed great diversity of molluscs during the one year study period. The Nakana Lake shows Shannon-Weiner Index (H) varied from 6.5137 to 3.0866 at Site-I and at Site –II in between 5.2641 to 2.2611. Simpson's Dominance Index (D) fluctuated in between 0.0887 and 0.1216 in Monsoon, 0.1183 and 0.1461 in Winter and 0.1212 and 0.1376 in Summer, where, Site-II was found to be dominant over Site-I. Values of Simpson's Index of Diversity at site –I is found to be varied i.e. 0.9113 in Monsoon, 0.8817 in Winter and 0.8788 in Summer Whereas, 0.8784 in Monsoon, 0.8539 in Winter and 0.8624 in Summer, at Site II. Simpson's reciprocal Index (1/D) was highest in Monsoon 11.273 and 8.2508 at Site-I and at Site-II and least in Winter 8.453 and 6.8446 respectively. Simpson's Evenness (E) maximum in Monsoon 0.8671 followed by Winter 0.6502 and Summer 0.6346 at Site-I and at site II maximum recorded in Monsoon 0.6853 followed by 0.6056 in Summer, 0.5703 in Winter. Pielou's Evenness (J) ranges in between 2.5395 to 2.1184 in Summer season at Site-I and Site-II respectively while 0.9099 to 1.2033 in Winter and

1.4867 to 1.2306 in monsoon. The present investigation resembles with the observation made by so many workers on the occurrence of molluscan diversity in different water bodies (Dollikar *et al.*, 2014, Mohan *et al.*, 2013).

CONCLUSION

In general, the present study revealed that species richness, abundance, density and diversity of molluscs depend upon rich ecosystem of Nakana Lake. All recorded molluscan species are indigenous. There found a considerable difference within the study localities.

ACKNOWLEDGEMENT

The authors are grateful to the Principal, S.G. Patil, A.C. S. College, Sakri, Dist. Dhule (M.S), India, for providing laboratory facilities during tenure of research work and also thankful to ZSI, Pune for scientific identification of Molluscan species.

REFERENCES

- Ahirrao KD. (2003).** Effect of sevin on some ecophysiological aspects of a Melanoid snail, *Melania scabra*. Ph. D. Thesis submitted to Dr. BAMU, Aurangabad.
- Ahirrao KD and Khedkar GD. (2012).** Variations in LC₅₀ of some pesticides due to physico-chemical parameters and impact of sevin on neurosecretory cells of a fresh water snail, *Thiara lineata*, *Journal of Research in Biology*, 1:050-056.
- El-kady GA, Shoukry A, Reda LA, El-Badri YS. (2000).** Survey and population dynamics of fresh water snail in newly settled areas of the Sinai Peninsula. *Egyptian Journal of Biology*, 2:42-48.
- Hussein MA, Obuid-Allah AH, Mahmoud AA, Fangary HM. (2011).** Population dynamics of fresh water snails (Mollusca:Gastropoda) at Qena Governorate, Upper Egypt. *Egypt. Acad. J. Biol. Sci.*, 3 (1):11-22.
- Hyman LH. (1967).** The invertebrates Vol.6. Mollusca, 1, McGraw Hill, New York.
- Micheal P. (1984).** Tata McGraw – Hill Publishing Company Limited, New Delhi.
- Patole VM. (2010).** Ecology and biodiversity of Mangroves in Mochamad estuary of Vengurla, South Konkan, Maharashtra. Ph. D. Thesis submitted to University of Mumbai.
- Pielou EC. (1975).** Ecological diversity. Wiley, New York,165.
- Sharma KK, Sharma SP and Sawhney N. (2009).** Distribution and ecology same freshwater molluscs of the Jammu Division of the J and K State. *J. Environ. Bio-sci.* 23(2):179-181.
- Sharma KK, Samita Chowdhary and Arti Sharma. (2010).** Malacofauna diversity of river Chenab fed stream(Gho-Manhasan), *Bioscan*, 6(2):267-279.
- Subba Rao NV. (1989).** Handbook: Freshwater molluscs of India. Zoological Survey of India, Calcutta.
- Dollikar AV, Mohite AS and Charde PN. (2014).** Correlation of molluscan diversity with physico-chemical characteristic of water of Gorewada reservoir, Nagpur, India. *International Journal of Life Sciences, Special issue*, A2:197-202.
- Mohan D, Elumalai V, Subbulakshmi G, Jayalakshmi S and Srinivasan M. (2013).** Biodiversity of cultivable molluscan resources from Pulicat Lake, Southeast coast of India, *Arthropods* 2(2): 53-65.
- Shannon EC and Wiener W. (1949).** The Mathematical theory of communication, University of Illinois Press Urbana, 125.
- Simpson EH. (1949).** Measurement of diversity, *Nature*,

163:688.

Preston HB. (1915). The fauna of British India including Ceylon and Burma. Mollusca (Freshwater Gastropoda and Pelecypoda). Taylor and Francis, Landon, i-xi+ 244.

Tonapi GT. (1980). Freshwater animals of india. An Ecological Approach. Oxford and IBH publishing co., New Delhi, Bombay, Calcutta 341.

Rao SNV. (1989): Freshwater Mollusca of India. In. Rao KS. (ED.). Recent advances in freshwater Biology. New Delhi. Anmol Publication. 2:187-202.

Submit your articles online at
www.jresearchbiology.com

Advantages

- **Easy online submission**
- **Complete Peer review**
- **Affordable Charges**
- **Quick processing**
- **Extensive indexing**
- **You retain your copyright**

submit@jresearchbiology.com

www.jresearchbiology.com/Submit.php