

Short Communication

Histopathology in the gut of the silkworm, *Bombyx mori* Linn- infected with microsporidia

Authors:

Shabir A. Bhat¹,
Ifat Bashir² and
Kamili AS³

Institution:

1. Temperate Sericulture
Research Institute (TSRI)
Mirgund, Sher-e-Kashmir
University of Agricultural
Sciences and Technology of
Kashmir. Post box No. 674,
G.P.O Srinagar-190001.

2. Basic Seed Station,
Mirgund, Sericulture
Development Department,
Kashmir, India

3. Directorate of Extension
Education, Sher-e-Kashmir
University of Agricultural
Sciences and Technology,
Kashmir, India.

Corresponding author:

Shabir A

ABSTRACT:

Ultrastructure of gut of the silkworm, *Bombyx mori* infected with microsporidia exhibited cytoplasmic vacuolization in the form of large empty spaces, fewer mitochondria, different spore stages (meronts and spronts) as grayish black spheres and mature spores. The meronts and spronts measured 0.61 and 0.56nm and 1.23 and 0.89nm in length and width respectively. The lightly infected gut, could not show any vacuolization but in the heavily infected gut cell cytoplasm destruction resulted in the formation of empty spaces.

Keywords:

Bombyx mori, Microsporidia, Mitochondria, Cytoplasm, Vacuolization.

Article Citation:

Shabir A. Bhat, Ifat Bashir and Kamili AS.

Histopathology in the gut of the silkworm, *Bombyx mori* Linn- infected with microsporidia

Journal of Research in Biology (2020) 10(6): 2872-2875

Dates:

Received: 10 Aug 2020 Accepted: 05 Sep 2020

Published: 28 Sep 2020

Web Address:

[http://jresearchbiology.com/
documents/RA0732.pdf](http://jresearchbiology.com/documents/RA0732.pdf)

This article is governed by the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which gives permission for unrestricted use, non-commercial, distribution and reproduction in all medium, provided the original work is properly cited.

INTRODUCTION

A number of microsporidia have been reported as infecting the gut of the silkworm and then passes to other organs *via* gut wall (Kawarabatta, 2003; Bhat, 2006). However, in most cases, reports have been dealt more with the development of the microsporidians with little emphasis on the host parasite interaction.

Microsporidia (Msp) isolated from economically important insect silkworm, *Bombyx mori* (Kamilli *et al.*, 2009 and 2011), infects the cytoplasmic cells of the gut causing histopathological alterations. This research describes the ultrastructural histopathological changes observed in microsporidian infected tissues of the

silkworm (Ananthalakshmi *et al.*, 1994).

MATERIALS AND METHODS

The 5th instar infected silkworms were dissected out dorsally under dissecting binocular microscope (Magnus-MS-24) and the gut was examined under binocular microscope (Olympus-CX31) as fresh mount. For histopathological studies, a portion of gut was cut and placed in fresh 3% (v/v) glutaraldehyde, fixed overnight at 4°C. The sample was post fixed in 1% osmium tetroxide for 2 h, washed, dehydrated in an ascending series of alcohol, passed through propylene oxide and infiltrated with araldite and propylene oxide for




Figure 1. Histopathology of infected gut of silkworm, a) arrows shows stage of infection b) multiplication of spores in cytoplasm c) arrows shows the rod shaped mitochondria d) meronts and sporonts and e) cytoplasm disorganization

12 h. The samples were centrifuged and sediment was infiltrated again with fresh araldite and kept at 60°C for 48 h. Ultra thin sections were double stained with uranyl acetate and lead citrate and was then observed and scanned under 60KVA (JEOL 100CX) electron microscope.

$$\text{Size of structure} = \frac{\text{Magnification of the reading}}{\text{Photo magnification}} \times \text{Scale unit selected (nm)}$$

The measurement of microsporidia with respect to different developmental stages was calculated by following the standard formula. The observations were recorded.

RESULTS AND DISCUSSION

The infected silkworms showed no external sign/ symptom. Hence they had to be dissected in order to determine their internal conditions. So the infection was diagnosed only from fresh smears of the gut. The ultrastructural studies showed presence of microsporidia spores and its various developmental stages in the infected tissues. Early developmental stages were intact with the gut epithelium and showed degeneration of cell organelles (Figure 1 a and b). Meronts, sporonts and mature spores could be observed in infected tissues. The meronts and sporonts measured 0.61, 0.56nm and 1.23, 0.89nm in length and width respectively. The length-width ration measured 1.09:1, 1.37:1 in meront and sporont respectively (Table 1). It was also observed that microsporidia multiplies in the cytoplasm of the gut cells but not in the nucleus, as the cytoplasm of the infected tissues was full of infection. Rod shaped mitochondria could also be observed between the gut epithelium (Figure 1 c). Cell degeneration due to large grayish spore developmental stages was obvious (Figure 1 d). Sporoblast in the early stages of the differentiation were seen in the infected cytoplasm and an another interesting observation was recorded that the infected cytoplasm

Table 1. Measurement of developmental stages of microsporidian, in the infected tissues of the mulberry silk worm

Developmental stages of the microsporidia	Size (nm)		
	Length	Width	Length-width ratio
Meront	0.61	0.56	1.09:1
Sporont	1.23	0.89	1.37:1

showed different developmental stages of the pathogen (Figure 1 e). Heavy infection load was observed in the form of different developmental spore stages and mature spores which resulted in marked cell disorganization, and empty spaces (without any cell organelle around the microsporidia).

The studies conducted on the development of microsporidian infection in silkworm showed chronic effect as the infected silkworm showed normal growth up to spinning and did not exhibit any visible sign of infection externally. Contrary to this study, Baig (1994) made observations on the silkworm infected with *Nosema bombycis* and found that it showed sluggish movement, tiny size, loss of appetite, vomit of gut juice and diarrhea with also pepper like spots on the body of the silkworms. The ultra structure studies of gut revealed that although the cytoplasmic organelles of lightly infected cells are relatively organized, however, marked disorganization occurred in the heavily infected cell cytoplasm. The microsporidia usually occur in the cytoplasm but there are few reports of their occasional occurrence in the cell nucleus such as *Nosema apis* in honey bees, *Chorestoneura fumifera* in the spruce and *N. bombycis* in the silkworm (Tanada and Kaya, 1993). It is in conformity with the results of Tanada and Kaya, (1993) that microsporidian (Msp) was also located in the cell cytoplasm. The present study revealed that the marked cytoplasmic destruction, reduction of cell organelles and occurrence of large empty spaces was due to the microsporidia infection. It is concluded that cell organelle destruction was due to the accumulation of parasite within the tissues and not the consequence of

any kind of active function performed by the pathogen. In a similar study, Jurad *et al.* (1967) reported the microsporidia infection caused marked ultra structure changes in the cells of the salivary gland of the Sciarid flies due to deletion, disorganization of cell organelles and vacuolation of the cytoplasm. Jyothi *et al.* (2005) reported the large empty spaces around the microsporidia without any cell organelle, the developing stages of the microsporidia spores which are larger in size and occupy enormous spaces and once it reaches its maturity, it shrunk in size and the spaces remain as such or in other words lysis of the host cytoplasm takes place.

ACKNOWLEDGEMENT

The authors are thankful to transmission electron microscopy unit of National Institute of Mental Health and Neuro Sciences (NIMHANS), Bangalore, for providing the laboratory facilities to carry out this work.

REFERENCES

Ananthalakshmi KVV, Fujiwara T and Datta RK. 1994. First report on the isolation of three microsporidians (*Nosema* spp.) from the silkworm, *Bombyx mori* L. in India. *Indian Journal of Sericulture*, 33: 146-48.

Baig M. 1994. Studies on *Nosema bombycis* - A pathogen of silkworm *Bombyx mori* L. PhD Thesis, University of Mysore, Mysore.

Bhat SA. 2006. Characterization of microsporidian infecting lamerin breed of the silkworm, *Bombyx mori* L. PhD Thesis. University of Mysore, Mysore.

Cavalier-Smith T. 1987. Eukaryotes with no mitochondria. *Nature*, 326: 332-333.

Jurand AL, Simoes LS and Pavan C. 1967. Changes in the ultra structure of salivary gland cytoplasm in *Sciara ocellaris* (Comstock, 1982) due to the microsporidia infection. *Journal of Insect Physiology*,

13: 795-803.

Jyothi NB, Patil CS and Dass CMS. 2005. Effect of *Nosema bombycis* Naegeli (Nosematidar) infection on the silk gland of silkworm *Bombyx mori* L (Bombycidae) – an ultrastructural study. *Sericologia*, 45 (4): 413-421.

Kawarabata T. 2003. Biology of microsporidians infecting the silkworm, *Bombyx mori*, in Japan. *Journal of Insect Biotechnology and Sericology*, 72(1): 1-32.

Kamilli SA, Bhat SA and Bashir I. 2011. Observations on the microsporidia isolated from bivoltine mulberry silkworm *Bombyx mori* L reared under temperate climatic conditions in India. In 22nd congress of the international sericultural commission, ‘silk for the better life’; 2011 December 14-18; Chiang Mai Thailand. 110-117 P.

Kamilli SA, Bhat SA, Malik MA, Sofi AM, Malik GN and Zargar MA. 2009. Studies on microsporidia isolated from silkworm, *Bombyx mori* L. In 9th agricultural science congress; 2009 June 22-24. University of agricultural sciences and technology of Kashmir, India. 19 P.

Tanada Y and Kaya KH. 1993. Insect pathology, Academic Press Inc., New York. 441-458 P.

Submit your articles online at www.jresearchbiology.com

Advantages

- Easy online submission
- Complete Peer review
- Affordable Charges
- Quick processing
- Extensive indexing
- You retain your copyright

submit@jresearchbiology.com

www.jresearchbiology.com/Submit.php